

## 9.a L'osteoporosi: la malattia

Chiara Cepollaro, Maria Luisa Brandi

- Adami S., Bertoldo F., Brandi M.L., Cepollaro C., Filippini P., Fiore E. et al. (2009), *Guidelines for the diagnosis, prevention and treatment of osteoporosis*, in «Reumatismo», 61, pp. 260-284.
- Adami S., Giannini S., Giorgino R., Isaia G., Maggi S., Sinigaglia L. et al. (2003), *The effect of age, weight, and lifestyle factors on calcaneal quantitative ultrasound: The ESOPO study*, in «Osteop. Int.», 14, pp. 198-207.
- Black D.M., Arden N.K., Palermo L., Pearson J., Cummings S.R. (1999), *Prevalent vertebral deformities predict hip fractures and new vertebral deformities but not wrist fractures*, in «J. Bone Miner. Res.», 14, pp. 821-828.
- Cauley J.A., Thompson D.E., Ensrud K.C., Scott J.C., Black D. (2000), *Risk of mortality following clinical fractures*, in «Osteop. Int.», 11, pp. 556-561.
- Center J.R., Nguyen T.V., Schneider D., Sambrook P.N., Eisman J.A. (1999), *Mortality after all major types of osteoporotic fracture in men and women: An observational study*, in «Lancet», 353, pp. 878-882.
- Cianferotti L., Brandi M.L. (2012), *Guidance for the diagnosis, prevention and therapy of osteoporosis in Italy*, in «Clin. Cas. Miner. Bone Metab.», 9, pp. 170-178.
- Consensus Development Conference (2001), *Osteoporosis: Prevention, diagnosis, and therapy*, in «JAMA», 285, pp. 785-795.
- Cooper C. (1997), *The crippling consequences of fractures and their impact on quality of life*, in «Am. J. Med.», 103(2A), pp. 12-19.
- Ernest E. (1998), *Exercise for female osteoporosis. A systematic review of randomized clinical trials*, in «Sport. Med.», 25, pp. 359-368.
- Gold D.T. (1996), *The clinical impact of vertebral fractures: Quality of life in women with osteoporosis*, in «Bone», 18, pp. 185-189.
- Kanis J.A., Johnell O., Oden A., Johansson H., McCloskey E. (2008), *FRAX and the assessment of fracture probability in men and women from the UK*, in «Osteop. Int.», 19, pp. 385-397.
- Kanis J.A. on behalf of the WHO Scientific Group (2008), *Assessment of osteoporosis at the primary health-care level*, technical report, WHO Collaborating Centre, University of Sheffield, Sheffield, [www.shef.ac.uk/FRAX](http://www.shef.ac.uk/FRAX).
- Lewiecki E.M., Kendler D.L., Kiebzak G.M., Schmeer P., Prince L.R., El-Hajj Fuleihan G. et al. (2004), *Special report on the official position of the International Society for Clinical Densitometry*, in «Osteop. Int.», 15, pp. 779-784.
- Lindsay R., Silverman S.L., Cooper C., Hanley D.A., Barton I., Broy S.B. (2001), *Risk of new vertebral fracture in the year following a fracture*, in «JAMA», 285(3), pp. 320-323.
- National Osteoporosis Foundation (1998), *Osteoporosis: Review of the evidence for prevention, diagnosis and treatment and cost-effectiveness analysis. Status report*, in «Osteop. Int.», 8, pp. 8-16.
- NIH Consensus Development Panel (1994), *Optimal calcium intake. NIH consensus conference*, in «JAMA», 272, pp. 1242-1248.
- Randall A.G., Nguyen T.V., Bhalerao N., Silverman S.L., Sambrook P.N., Eisman J.A. (2000), *Deterioration in quality of life following hip fracture: A prospective study*, in «Osteop. Int.», 11, pp. 460-466.

Rizzoli R., Branco J., Brandi M.L., Boonen S., Bruyère O., Cacoub P. et al. (2014), *Management of osteoporosis of the oldest old*, in «Osteop. Int.», 25(11), pp. 2507-2529.

Van der Wielen R.P., Lowik M.R., Van den Berg H., De Groot L.C., Haller J., Moreiras O. et al. (1995), *Serum vitamin D concentrations among elderly people in Europe*, in «Lancet», 346, pp. 207-210.

WHO Study Group (1994), *Assessment of fracture risk and its application to screening for postmenopausal osteoporosis: Report of a WHO Study Group*, technical report series 843, WHO, Genève.